arXiv:1206.2387v2 [math.GT] 18 Nov 2012

THE DEFINABILITY CRITERION FOR COCOMPACT CONVEX PROJECTIVE POLYHEDRAL REFLECTION GROUPS

KANGHYUN CHOI AND SUHYOUNG CHOI

Abstract. In this paper, we prove the criterion for a Zariski dense subgroup generated by reflections $\Gamma \subset \mathrm{SL}^{\pm}(n+1,\mathbb{R})$ to be definable over $\mathbb{A}$ where $\mathbb{A}$ is an integrally closed Noetherian ring in the field $\mathbb{R}$. We apply this criterion for groups generated by reflections that act cocompactly on irreducible properly convex open subdomains of the n-dimensional projective sphere. This gives a methodology to construct injective group homomorphisms from such Coxeter groups to $\mathrm{SL}^{\pm}(n+1,\mathbb{Z})$. Finally we provide some examples of $\mathrm{SL}^{\pm}(n+1,\mathbb{Z})$-representations of such Coxeter groups. In particular, we consider simplicial reflection groups that are isomorphic to hyperbolic simplicial groups and classify all the conjugacy classes of the reflection subgroups in $\mathrm{SL}^{\pm}(n+1,\mathbb{R})$ that are definable over $\mathbb{Z}$.

1. Introduction

Let $\mathbb{F}$ be a field of characteristic 0 and let V be an (n+1)-dimensional vector space over $\mathbb{F}$. We define

$$\mathrm{SL}^{\pm}(n+1,\mathbb{F}) := \{A \in \mathrm{GL}(n+1,\mathbb{F}) | \det A = \pm 1\}$$

as the group of linear transformations from V to V whose determinants are either 1 or -1. A reflection σ is an element of order 2 of $\mathrm{SL}^{\pm}(n+1,\mathbb{F})$ which is the identity on an hyperplane. Every reflection σ acts on V as $\sigma_{\alpha,\vec{v}} : \vec{x} \to \vec{x} - \alpha(\vec{x})\vec{v}$. Namely, all reflections are of the form $\sigma := \sigma_{\alpha,\vec{v}} := I - \alpha \otimes \vec{v}$ for some $\alpha \in V^*$ and $\vec{v} \in V$ with $\alpha(\vec{v}) = 2$ where $\vec{v}$ and α are different from zero and determined within a transformation $\vec{v} \to c\vec{v}$, $\alpha \to c^{-1}\alpha$. Let Γ be the semigroups generated by reflections $\sigma_i = \sigma_{\alpha_i,\vec{v}_i}$, $i \in \mathbb{N}$. We set $c_{i,j} = \alpha_i(\vec{v}_j)$ and call the (possibly infinite) matrix $C = (c_{i,j})$ the *Cartan matrix* of Γ. Products of form $c_{i_1 i_2} c_{i_2 i_3} ... c_{i_k i_1}$ are called *cyclic products*. A cyclic product is called *simple* if all its indices $i_1, ..., i_k$ are distinct. It is clear that every cyclic product is a product of simple ones.

Let O denote the origin in $\mathbb{R}^{n+1}$ from now on. Let $\mathbb{S}^n$ be the n-dimensional real projective sphere, i.e. the quotient $\mathbb{R}^{n+1} - \{O\}/\sim$ where $\sim$ is the equivalence relation given by that $\vec{v} \sim \vec{w}$ if and only if $\vec{v} = r\vec{w}$ for a positive real number r. Then we can identify $\mathrm{SL}^{\pm}(n+1,\mathbb{R})$ with the group of projective transformations of $\mathbb{S}^n$. Denote by $\Pi : \mathbb{R}^{n+1} - \{O\} \to \mathbb{S}^n$ the quotient map.

Vinberg [28] initiated a systematic study to construct injective group homomorphisms from a Coxeter group that is geometrically realized as an n-dimensional polyhedron to the group $\mathrm{SL}^{\pm}(n+1,\mathbb{R})$: the groups generated by reflections by sides of the n-dimensional polyhedron in the n-dimensional real projective sphere $\mathbb{S}^n$. Based on Vinberg's results, the properties of the corresponding representation

---

Both authors were supported by Mid-career Researcher Program through NRF grant (2011-0027600) funded by the MEST..

spaces are studied by Benoist([2], [3], [4], [5]), Choi([10]), and Marquis([21]). See proposition 6.2 and figures1-4 in section 6 for the parametrized subspace in the case of triangular reflection group.

A *great circle* in $\mathbb{S}^n$ is the image $\Pi(V - \{O\})$ for a two-dimensional subspace V in $\mathbb{R}^{n+1}$. A *segment* is a proper connected subset of the great circle. A *convex* segment is one that does not contain an antipodal pair of points except at the end points. A *closed hemisphere* is the image $\Pi(H - \{O\}$ where H is a subset of $\mathbb{R}^{n+1}$ given by a linear inequality $f(\vec{v}) \geq 0$. An *open hemisphere* is the interior of a closed hemisphere. A *convex subset* of $\mathbb{S}^n$ is a set such that any pair of its point is connected by a convex segment. A convex subset is either $\mathbb{S}^n$ itself or a subset of a closed hemisphere given by a linear inequality. A *properly convex subset* is a convex subset whose closure does not contain a pair of antipodal points. It is a precompact subset of an open hemisphere H.

Now suppose that the n-dimensional properly convex polyhedron P is compact in $\mathbb{S}^n$. Let Γ be the group generated by reflections by sides of P satisfying edge order conditions (the conditions in Proposition 3.1). Let Δ be a family of linear transformations of V. A ring $\mathbb{A}$ in $\mathbb{F}$ is called a *ring of definition* for Δ if V contains an $\mathbb{A}$-lattice that is invariant under Δ. In that case we also say that Δ is *definable over* $\mathbb{A}$. If $\mathbb{A}$ is a principal ideal domain, then the fact that Δ is definable over $\mathbb{A}$ is equivalent to that there is a basis in which the transformations of Δ can be written down by matrices with elements in $\mathbb{A}$. Our main theorem is Theorem 1.1. We defer precise definitions in sections 2, 3, and 4.

Theorem 1.1. *Suppose* $n \geq 2$ *be an integer. Let* $\Gamma \subset \mathrm{SL}^{\pm}(n+1, \mathbb{R})$ *be a group generated by reflections that divides an irreducible strictly convex open domain* Ω *in* $\mathbb{S}^n$*. Let* Δ *be any subgroup of finite index in* Γ*. Then the following properties of an integrally closed Noetherian ring* $A \subset \mathbb{R}$ *are equivalent.*

(1) Γ *is definable over* $\mathbb{A}$*.*
(1′) $\mathbb{A}$ *contains all the simple cyclic products of the Cartan matrix of* Γ*.*
(2) Δ *is definable over* $\mathbb{A}$*.*

See Section 6 and Proposition 6.2 for concrete examples of such group actions.

Our strategy to prove Theorem 1.1 is to prove that the theorem holds for Zariski dense reflection groups in $\mathrm{SL}^{\pm}(n+1, \mathbb{F})$ where $\mathbb{F}$ is an algebraically closed field of characteristic 0. This result is Theorem 5.3. In the proof, Vinberg's theory on ring of definitions of Zariski dense subgroups of linear algebraic group is used. In particular, we made use of the fact that if Δ is definable over $\mathbb{A}$ then $\mathrm{Ad}(\Gamma)$ is definable over $\mathbb{A}$. With Vinberg's theory, Lemma 5.2 is essential for the proof. Theorem 5.3, Vinberg's result for orthognal groups, and Benoist's Zariski density theorem (Theorem 2.1) imply Theorem 1.1. Our example of polyhedral reflection groups that are definable over $\mathbb{Z}$ includes some triangular, tetrahedral, and cubical reflection groups. Indeed the motivation for our theoretical results come from Vinberg and Kac's example of a $(3, 3, 4)$-triangle reflection group in $\mathrm{SL}^{\pm}(3, \mathbb{Z})$ ([26]).

The equivalence of (1) and (2) in Theorem 1.1 can be proved even for Zariski dense subgroups generated by finitely many elements of finite order if we add some conditions. This result is Theorem 5.3 which we state here.

Theorem 1.2. *Suppose* $n \geq 2$ *is an integer. and let* $\mathbb{F}$ *be an algebraically closed field of characteristic* 0*. Let* $\Gamma \subset \mathrm{SL}^{\pm}(n+1, \mathbb{F})$ *be a Zariski dense subgroup generated by finitely many elements* σ_i*s for* $i = 1, \cdots, m$*. Let* Δ *be any subgroup of finite index* *in* Γ. *Let* $\mathbb{A}$ *be an integrally closed Noetherian ring in* $\mathbb{F}$, *and let* $\mathbb{K}$ *be the field of fractions of* $\mathbb{A}$. *Suppose that for* $i = 1, \cdots, m$, σ_i *satisfies one of the following conditions*

- *The order of* σ_i *is finite and relatively prime to* $n+1$.
- *The trace of* σ_i *is nonzero and in* $\mathbb{K}$.

Then the following properties of an integrally closed Noetherian ring $\mathbb{A}$ *in* $\mathbb{F}$ *are equivalent.*

(1) Γ *is definable over* $\mathbb{A}$.
(2) Δ *is definable over* $\mathbb{A}$.

Section 2 reviews basic definitions of real projective orbifolds and state Benoist's Zariski density theorem. Section 3 reviews Vinberg's theory of polyhedral reflection groups. Section 4 reviews Vinberg's theory on rings of definition of Zariski dense subgroups of semisimple linear algebraic groups. Section 5 states and prove our main results. Finally, in Section 6 we apply our main results for some convex projective polyhedral reflection groups to determine if they are definable over $\mathbb{Z}$. In particular, we will consider convex projective simplicial reflection groups that are isomorphic to hyperbolic simplicial groups and classify all the conjugacy classes of reflection groups that are definable over $\mathbb{Z}$. (See Theorem 6.2, 6.5, and 6.6.)

Our results are applicable to many hyperbolic Coxeter orbifolds including orderable ones [10] and Marquis's [21] and we plan to study these in later papers.

There are some people whose help and encouragement we would like to acknowledge. We thank William Goldman for introducing us Vinberg and Kac's example of a $(3,3,4)$-triangle reflection group. In fact, he has found some of these examples himself. We thank Alan Reid and Dave Witte Morris for giving us many informations on arithemetic groups, and Gye-Seon Lee for helpful discussions and for letting us use his graphics of a cube and a prism. We also thank Ja Kyung Koo and his students for having joint seminars on arithmetic hyperbolic geometry which helped us much.

2. Real projective orbifolds

2.1. **Orbifolds.** Given two manifolds M_1 and M_2 with groups G_1 and G_2 acting on them respectively, a map $f : M_1 \to M_2$ is *equivariant* with respect to a homomorphism $h : G_1 \to G_2$ if we have $h(g) \circ f = f \circ g$ for each $g \in G_1$.

We will be using the language of orbifolds for later purposes. An n-dimensional *orbifold structure* on a second countable Hausdorff space X is given by atlas of compatible charts (U, G, ϕ) where U is an open subset of $\mathbb{R}^n$, G is a finite group acting on U, and ϕ is a homeomorphism $\phi : U/G \to V$ for an open subset V of X. We say that $\phi(U/G)$ is *modeled* on (U, G) or (U, G, ϕ) also. Two charts (U, G, ϕ) and (U', G', ϕ') are *compatible* if given any point $p \in \phi(U/G) \cap \phi(U'/G')$, there is a chart (U'', G'', ϕ'') so that $\phi''(U''/G'')$ is an open neighborhood in $\phi(U/G) \cap \phi(U'/G')$ where there are smooth lifts $\tilde{\phi}''_1 : U'' \to U$ equivariant with respect to an imbedding $G'' \to G$ and $\tilde{\phi}''_2 : U'' \to U'$ equivariant with respect to $G'' \to G'$.

Here, X with an orbifold structure is said to be an *orbifold* and X is said to be the underlying space of the orbifold.

Given a manifold M and a discrete group Γ acting on M properly discontinuosly and but may be not freely, we can form M/Γ as a quotient space. M/Γ has a natural orbifold structure given by covering M by open sets U so that if $g(U) \cap U \neq \emptyset$, then

$g(U) = U$. The collection of all (U, G, ϕ) where G is the subgroup of Γ acting on an open set U gives us an orbifold structure.

Since the orbifolds we study here admit real projective structures, they are of form M/Γ for a manifold M and a discrete group Γ acting on M properly discontinuously and but may be not freely by Thurston [24]. Two such orbifolds M/Γ and N/Δ for simply connected M and N are *diffeomorphic* if there is a diffeomorphism $f : M \to N$ equivariant with respect to an isomorphism $h : \Gamma \to \Delta$. The *orbifold fundamental group* $\pi_1(Q)$ is the abstract group which is isomorphic to Γ whenever M is simply connected. Clearly, it acts on M. If M is simply connected, M is said to be a *universal cover* of M/Γ. (For the general definition of orbifold and geometric structures on it, we refer to [9] and [11].)

2.2. **Real projective orbifolds.** Let $n \geq 2$ and V be an n-dimensional real vector space $\mathbb{R}^{n+1}$. We identify

$$\mathrm{SL}^{\pm}(n+1, \mathbb{R}) := \{A \in \mathrm{GL}(n+1, \mathbb{R}) | \det A = \pm 1\}$$

with the group of projective transformations of $\mathbb{S}^n$. Recall that a subdomain Ω of $\mathbb{S}^n$ is *properly convex* if it is convex and its closure $\overline{\Omega}$ does not contain two opposite points. Ω is *strictly convex* if moreover its boundary $\partial\Omega$ does not contain any line segment of positive length.

A *properly convex projective orbifold*, Q, is of the form Ω/Γ where Ω is a properly convex open domain in $\mathbb{S}^n$ and Γ is a discrete subgroup of $\mathrm{SL}^{\pm}(n+1, \mathbb{R})$ which preserves Ω acting properly discontinuously. A *properly convex projective structure* on an orbifold M is a diffeomorphism $f : M \to Q$ for Q as above.

The two properly convex projective orbifolds $Q_1 = \Omega_1/\Gamma_1$ and $Q_2 = \Omega_2/\Gamma_2$ are *projectively diffeomorphic* if there exists a projective transformation $h \in \mathrm{SL}^{\pm}(n+1, \mathbb{R})$ such that $h(\Omega_1) = \Omega_2$ and $h\Gamma_1 h^{-1} = \Gamma_2$. When the properly convex projective orbifold $Q = \Omega/\Gamma$ is compact, we say that Γ *divides* Ω.

Given a properly convex projective structure on an orbifold M, we obtain a homomorphism $h : \pi_1(M) \to \mathrm{SL}^{\pm}(n+1, \mathbb{R})$, called a *holonomy homomorphism* determined up to conjugation by $\mathrm{SL}^{\pm}(n+1, \mathbb{R})$: Let Ω/Γ be a properly convex projective orbifold and $f : M \to \Omega/\Gamma$ be a projective diffeomorphism. Then let $f_* : \pi_1(M) \to \Gamma$ be the induced homomorphism. Then f_* considered as a map $\pi_1(M) \to \mathrm{SL}^{\pm}(n+1, \mathbb{R})$ is the holonomy homomorphism. The image of the holonomy homomorphism is called the *holonomy group* of M or $\pi_1(M)$.

Any open domain Ω in $\mathbb{S}^n$ is the image of the unique convex open cone C in V under the projection map. The open convex cone C in $V - \{O\}$ is said to be *reducible* if it can be written as the sum $C = C_1 \oplus C_2$ of two convex cones C_i in proper subspaces V_i of V. A properly convex open domain Ω in $\mathbb{S}^n$ is said to be *reducible* if its preimage C is reducible. Otherwise we say that Ω is *irreducible.* The *hyperbolic n-space* $\mathbb{H}^n$ is the space of rays in the Lorentz cone

$$\Lambda_{n+1} := \{x \in \mathbb{R}^{n+1} | q(x) > 0 \text{ and } x_1 > 0\}$$

where $q(x) = x_1^2 - x_2^2 - \cdots - x_{n+1}^2$. Its group of automorphisms is the group $\mathrm{Aut}(\mathbb{H}^n) = \mathrm{O}^+(1, n)$ of orthogonal transformations of q which preserve $\mathbb{H}^n$. If there exists a discrete subgroup Γ of $\mathrm{O}^+(1, n)$ which preserves $\mathbb{H}^n$, we call the quotient $\mathbb{H}^n/\Gamma$ a *hyperbolic orbifold.* $\mathbb{H}^n$ is an example of irreducible strictly convex open domains in $\mathbb{S}^n$. A group Γ of $\mathrm{SL}^{\pm}(n+1, \mathbb{R})$ is said to be *irreducible* if there are no

Γ-invariant nontrivial subspaces in $\mathbb{R}^{n+1}$, and is said to be *strongly irreducible* if all of its finite index subgroups are irreducible.

2.3. **Benoist's theorem.** Now we state version of Cooper and Delp (Corollary 4.2 of [15]) of Benoist's Zariski density theorem [1] .

Theorem 2.1. *Let* Γ *be a discrete subgroup of* $\mathrm{SL}^{\pm}(n+1,\mathbb{R})$ *which divides an irreducible strictly convex open domain* Ω *in* $\mathbb{S}^n$*. The real Zariski closure* $\overline{\Gamma}$ *is either* $\mathrm{O}^{+}(1,n)$ *or* $\mathrm{SO}^{+}(1,n)$ *iff* Ω *is* $\mathbb{H}^n$*. Otherwise* $\overline{\Gamma}$ *is either* $\mathrm{SL}^{\pm}(n+1,\mathbb{R})$ *or* $\mathrm{SL}(n+1,\mathbb{R})$*.*

Definition 2.2. A discrete subgroup of $\mathrm{SL}^{\pm}(n+1,\mathbb{R})$ which divides an irreducible properly convex open domain Ω in $\mathbb{S}^n$ is said to be *irreducible convex dividing linear group.*

Of course this group is irreducible subgroup of $\mathrm{SL}^{\pm}(n+1,\mathbb{R})$ by Vey irreducibility theorem (Theorem 5.1 of [6]).

3. Polyhedral reflection groups

3.1. **Vinberg's Condition.** Let $P \subset \mathbb{S}^n$ be a *n-dimensional convex polyhedron*, i.e. the radial image in $\mathbb{S}^n$ of a convex polyhedral cone of $\mathbb{R}^{n+1}$ with O omitted. A *k-face* of P is a k-dimensional convex subset of P obtained as an intersection of P with some hyperspheres which do not meet the interior P^o. A *face* is an $(n-1)$-face. A *projective reflection* is a reflection defined in n-dimensional real vector space, i.e. a reflection in $\mathrm{SL}^{\pm}(n+1,\mathbb{R})$. Let S be the set of faces of P and for every s in S, we can associate a projective reflection $\sigma_s = I - \alpha_s \otimes \vec{v}_s$ with $\alpha_s(\vec{v}_s) = 2$ which fixes s. A suitable choice of signs allows us to suppose that P is defined by the inequalities $(\alpha_s \leq 0)_{s\in S}$. If $s \cap t$ is nonempty and θ is the dihedral angle, then $\sigma_s\sigma_t$ is the rotation around the 2-codimensional face $s \cap t$ by 2θ. Let $c_{s,t} := \alpha_s(\vec{v}_t)$ for $s, t \in S$. Let Γ be the group generated by the reflections σ_s. The following theorem of Vinberg [28] provides a necessary and sufficient condition for such a group Γ acts on a convex subdomain of $\mathbb{S}^n$. Note here that $c_{s,t}c_{t,s} = 4\cos^2(\frac{\pi}{m_{s,t}})$ is the condition that $\sigma_s\sigma_t$ generate a cyclic group of order $m_{s,t}$.

Proposition 3.1. *Let* P *be a convex polyhedron of* $\mathbb{S}^n$ *and, for each face* s *of* P*, let* $\sigma_s = I - \alpha_s \otimes \vec{v}_s$ *be a projective reflection fixing this face* s*. Let* Γ *be the group generated by the reflections* σ_s*. Then the following conditions for every* s, t *such that codim*$(s \cap t) = 2$ *are necessary and sufficient in order for* Γ *to preserves some convex subdomain* Ω *of* $\mathbb{S}^n$ *with the fundamental domain* P

(1) $c_{s,t} \leq 0$ *and* $(c_{s,t} = 0 \Leftrightarrow c_{t,s} = 0)$
(2) $c_{s,t}c_{t,s} \geq 4$ *or* $c_{s,t}, c_{t,s} = 4\cos^2(\frac{\pi}{m_{s,t}})$ *with* $m_{s,t} \geq 2$ *integer*

Moreover the group Γ *is discrete. The convex domain* Ω *is open if and only if for every* x *in* P*, the group* Γ_x *generated by* σ_s *for* s *containing* x *is a finite group. In this case,* Γ *acts properly on* Ω *with compact quotient.*

We will call the group generated by projective reflections fixing sides of some n-dimensional convex polyhedron P of $\mathbb{S}^n$ the *polyhedral reflection group*. Any polyhedral reflection group has the presentation

$$\langle s_i | (s_i s_j)^{n_{ij}} = 1\rangle,\ n_{ij} \in \mathbb{N},\ i, j = 1, \ldots, n,$$

where n_{ij} is defined for a subset of $\{1, \dots, n\}^2$ and n_{ij} is symmetric in i, j and $n_{ii} = 2$. An abstract group which has the above presentation is called *Coxeter group* with n generators. When the polyhedral reflection group Γ preserves a properly convex domain Ω, it is said to be *convex projective* and we call the quotient orbifold Ω/Γ of an open manifold Ω a *convex projective polyhedral reflection orbifold.*

3.2. **Coxeter orbifold.** A *reflection* in an open subset U of $\mathbf{R}^n$ is a transformation $U \to U$ of order two fixing a hypersurface in it. An *n-dimensional Coxeter orbifold structure* on an n-dimensional polyhedron P is given by giving orbifold structure on P where each point of the interior of each face has a chart modeled on an open subset in $\mathbf{R}^n$ with a reflection acting on it and each point of the interior of each side of codimension 2 is has a chart modeled on an open subset of $\mathbf{R}^n$ with a dihedral group generated by reflections. The polyhedron P with the Coxeter orbifold structure is denoted by $\hat{P}$ and is said to be an n-dimensional Coxeter orbifold. If P is compact, then $\hat{P}$ is said to be *compact*. In our cases, $\hat{P} = M/\Gamma$ for a simply-connected manifold M and Γ is a discrete group acting properly discontinuously since $\hat{P}$ admits a real projective structure. (See [9] for details.)

If M is a convex domain in $\mathbb{S}^n$ and a polyhedral reflection group Γ acts cocompactly and properly discontinuously, we say that Γ is a *convex projective reflection group*. In this case, the orbifold M/Γ has *convex projective structure*. If Γ acts on $\mathbb{H}^n$, then Γ is a *hyperbolic reflection group*. In the later case, M/Γ has *hyperbolic structure*.

Proposition 3.2.
- *Given two compact Coxeter orbifolds M_1 and M_2, M_1 is diffeomorphic to M_2 if and only if $\pi_1(M_1)$ is isomorphic to $\pi_1(M_2)$.*
- *The image of the homomorphism $\pi_1(M_1) \to \mathrm{SL}^{\pm}(n+1, \mathbb{R})$ divides a properly convex domain in $\mathbb{S}^n$ if and only if it is a holonomy homomorphism of a compact properly convex projective Coxeter orbifold M_1.*
- *the set of cocompact properly convex irreducible holonomy homomorphisms of M_1) is identical with the set of irreducible homomorphisms of $\pi_1(M_1) \to \mathrm{SL}^{\pm}(n+1, \mathbb{R})$ dividing a properly convex domains.*
- *This is also true if both sets are quotiented by the conjugation action by $\mathrm{SL}^{\pm}(n, \mathbb{R})$.*

Proof. Let P_1 and P_2 be the polyhedrons that are the respective underlying spaces of M_1 and M_2. By Example 7.14 of [13], these are $\pi_1(M_1)$ and $\pi_1(M_2)$ are groups of type PM, If $\pi_1(M_1)$ is isomorphic to $\pi_1(M_2)$, then the combinatorial data of the faces of P_1 and P_2 are identical by Theorem 13.4.1 of [13]. Hence, there exists a homeomorphism $M_1 \to M_2$ and $P_1 \to P_2$. A *stratum* of a Coxeter orbifold is one where the local groups of points are locally conjugate to one another by isotopies. (See Section 4.5 of [11] for definitions of strata, tubular neighborhoods, and so on.) Each stratum of an orbifold is diffeomorphic to a cell. We can smooth the map from lowest dimension stratum to maximal dimensional ones by induction and since a diffeomorphism defined on an open neighborhood of the boundary of a cell always extends to a diffeomorphism defined on a neighborhood of a cell. The first item follows.

For the second item, suppose that Ω be the properly convex domain and Γ be the image of the homomorphism k from $\pi_1(M)$ dividing Ω. Then Ω/Γ has the fundamental group isomorphic to Γ isomorphic to $\pi_1(M)$ and hence Ω/Γ is diffeomorphic to M_1 by a diffeomorphism $f : M_1 \to \Omega/\Gamma$ so that $f_* = k$. Therefore, k is a holonomy homomorphism of a properly convex projective structure on M_1.

Conversely, if k is a holonomy homomorphism of such a structure on M_1, then $k(\pi_1(M_1))$ acts on a properly convex domain Ω dividing it by definition.

The rest of these are consequences of the two items. $\square$

4. Vinberg's results on rings of definitions

In this section we state main results of Vinberg's paper [29] regarding rings of definition of Zariski dense subgroups of semisimple linear algebraic groups. Our main result in the next section is heavily based on Vinberg's theory..

Let V be a $(n+1)$-dimensional vector space over field $\mathbb{F}$ and $\mathbb{A}$ be a subring of $\mathbb{F}$. A set L in V is called an $\mathbb{A}$-*lattice* if it is a finitely generated A-submodule and the natural map $\mathbb{F} \bigotimes_{\mathbb{A}} L \to V$ is an isomorphism. If $\mathbb{A}$ is a principal ideal domain, then every $\mathbb{A}$-lattice has a basis which at the same time is a basis of V over $\mathbb{F}$. If $\mathbb{K}$is a subfield of $\mathbb{F}$, then a $\mathbb{K}$-lattice is just a $(n+1)$-dimensional vector space over $\mathbb{K}$.

Let Δ be a family of linear transformations of V. An integrally closed Noetherian ring $\mathbb{A}$ in $\mathbb{F}$ is called a *ring of definition* for Δ if V contains an $\mathbb{A}$-lattice that is invariant under Δ. In that case we also say that Δ is *definable over* $\mathbb{A}$. If $\mathbb{A}$ is a principal ideal domain, then the fact that Δ is definable over $\mathbb{A}$ means that there is a basis in which the transformations of Δ can be written down by matrices with elements in $\mathbb{A}$. If Δ is definable over $\mathbb{A}$ and $\mathbb{B}$ is a ring that contains $\mathbb{A}$, then Δ is also definable over $\mathbb{B}$.

Now we assume that $\mathbb{F}$ is an algebraically closed field of characteristic 0, G a semisimple algebraic group over $\mathbb{F}$ (not necessarily connected), and Γ a Zariski dense subgroup. Let Ad be the adjoint representation of G. Then Ad is a map from G to the automorphism group of its lie algebra $\mathfrak{g}$. For any matrix group H, let $\mathrm{tr}H$ be the set $\{\mathrm{tr}h | h \in H\}$. Following propositions will be useful in later sections.

Proposition 4.1 (Theorem 1 of [29]). *An integrally closed Noetherian ring* $\mathbb{A}$ *in* $\mathbb{F}$ *is a ring of definition for* $\mathrm{Ad}(\Gamma)$ *if and only if* $\mathbb{A} \supset \mathrm{tr}\mathrm{Ad}(\lambda)$ *for all* $\lambda \in \Gamma$.

Corollary 4.2 (Corollary to Theorem 1 of [29]). *There exists a smallest field of definition for the group* $\mathrm{Ad}(\Gamma)$. *If this field is an algebraic number field, then there exists a smallest ring of definition for this group.*

Proposition 4.3 (Theorem 2 of [29]). *Let* $\mathbb{A}$ *be an integrally closed Noetherian ring in* $\mathbb{F}$ *and* $\mathbb{K}$ *its field of fractions. Then* Γ *is definable over* $\mathbb{A}$ *if and only if it is definable over* $\mathbb{K}$ *and* $\mathrm{Ad}(\Gamma)$ *is definable over* $\mathbb{A}$.

Proposition 4.4 (Theorem 3 of [29]). *Let* Γ_1 *be a finite index subgroup of* Γ. *Then the classes of rings of definition for* $\mathrm{Ad}(\Gamma_1)$ *and* $\mathrm{Ad}(\Gamma)$ *are identical.*

Let V be an $(n+1)$-dimensional vector space over a (not necessarily algebraically closed) field $\mathbb{F}$ and V^* it dual. Recall that a linear transformation σ of V is called $reflection$ if it is an element of order 2 of $\mathrm{SL}^{\pm}(n+1, \mathbb{F})$ which is the identity on a hyperplane. Let Γ be a group generated by reflections. We denote by $\mathbb{Z}[\mathrm{tr}\Gamma]$ the ring with unit element generated by the set $\mathrm{tr}\Gamma$ in $\mathbb{F}$.

Proposition 4.5 (Lemma 11 of [29]). $\mathbb{Z}[\mathrm{tr}\Gamma]$ *is the ring with unit element generated by the simple cyclic products.*

A group H in $\mathrm{SL}^{\pm}(n+1,\mathbb{F})$ is *absolutely irreducible if* if it is irreducible as a matrix group in $\mathrm{SL}^{\pm}(n+1,\bar{\mathbb{F}})$ where $\bar{\mathbb{F}}$ is the algebraic closure of $\mathbb{F}$.

Proposition 4.6 (Lemma 12 of [29])**.** *Suppose that char$\mathbb{F} = 0$ and Γ is absolutely irreducible. An integrally closed Noetherian ring $\mathbb{A}$ in $\mathbb{F}$ is a ring of definition for Γ if and only if $\mathbb{A}$ contains $\mathbb{Z}[\mathrm{tr}\Gamma]$.*

Suppose that a nondegenerate scalar product $(,)$ is defined in V. The group of automorphisms of V that preserve this scalar product $(,)$ is denoted by $\mathrm{O}(V)$ and its unimodular subgroup by $\mathrm{SO}(V)$.

Proposition 4.7 (Theorem 5 of [29])**.** *Suppose $n \geq 3$, and let $\mathbb{F}$ be a algebraically closed field of characteristic zero. Let $\Gamma \subset \mathrm{O}(V)$ be a Zariski dense subgroup generated by reflections. Let Δ be any subgroup of finite index in Γ. Then the following properties of an integrally closed Noetherian ring $\mathbb{A} \subset \mathbb{F}$ are equivalent.*

(1) *Γ is definable over $\mathbb{A}$.*
(1′) *$\mathbb{A}$ contains all the simple cyclic products of the Cartan matrix of Γ.*
(2) *Δ is definable over $\mathbb{A}$.*
(3) *$\mathrm{Ad}(\Delta)$ is definable over $\mathbb{A}$.*
(4) *$\mathrm{Ad}(\Gamma)$ is definable over $\mathbb{A}$.*

5. The main result

Theorem 1.1 will be proved in this section. Throughout this section, let $\mathbb{F}$ be an algebraically closed field of characteristic zero. To prove Theorem 1.1, we prove that the similar result holds for finitely generated groups if we add some conditions.

5.1. The main result.

Theorem 5.1. *Suppose $n \geq 2$ is an integer. and let $\mathbb{F}$ be an algebraically closed field of characteristic 0. Let $\Gamma \subset \mathrm{SL}^{\pm}(n+1,\mathbb{F})$ be a Zariski dense subgroup generated by finitely many elements σ_is for $i = 1, \cdots, m$. Let Δ be any subgroup of finite index in Γ. Let $\mathbb{A}$ be an integrally closed Noetherian ring in $\mathbb{F}$, and let $\mathbb{K}$ be the field of fractions of $\mathbb{A}$. Suppose that for $i = 1, \cdots, m$, σ_i satisfies one of the following conditions*

- *The order of σ_i is finite and relatively prime to $n+1$.*
- *The trace of σ_i is nonzero and in $\mathbb{K}$.*

Then the following properties of an integrally closed Noetherian ring $\mathbb{A}$ in $\mathbb{F}$ are equivalent.

(1) *Γ is definable over $\mathbb{A}$.*
(2) *Δ is definable over $\mathbb{A}$.*

We consider the Lie algebra $\mathfrak{sl}_{n+1}(\mathbb{R})$ as a subalgebra of the matrix algebra $\mathbf{M}_n(\mathbb{R})$ which is realized from the decomposition $\mathbf{M}_{n+1}(\mathbb{R}) = \mathfrak{sl}_{n+1}(\mathbb{R}) + \mathbb{R} \cdot I_{n+1}$.

Before we prove Theorem 5.1, we need the following lemma of Grinberg which is available from http://mathoverflow.net/questions/81855/adjoint-map-and-number-field.

Lemma 5.2. *Let $\mathbb{L}$ be a field of characteristic 0, and $\mathbb{K}$ be a subfield of $\mathbb{L}$. Let $n \in \mathbb{N}$. Let $U \in \mathrm{GL}(n,\mathbb{L})$ be such that every $V \in \mathfrak{sl}_n(\mathbb{K})$ satisfies $UVU^{-1} \in \mathbf{M}_n(\mathbb{K})$. Then, there exists a nonzero $\lambda \in \mathbb{L}$ such that $\lambda U \in \mathrm{GL}(n,\mathbb{K})$.*

Proof. First, let us show that

(1) every $V \in \mathbf{M}_n(\mathbb{K})$ satisfies $UVU^{-1} \in \mathbf{M}_n(\mathbb{K})$.

Since $\mathbb{K}$ has characteristic 0, we have $\mathbf{M}_n(\mathbb{K}) = \mathfrak{sl}_n(\mathbb{K}) + \mathbb{K} \cdot I_n$, where I_n is the n by n identity matrix. Since (1) is linear in V, we are therefore done with the proof of (1) once we have shown that (1) holds for all $V \in \mathfrak{sl}_n(\mathbb{K})$ and for all $V \in \mathbb{K} \cdot I_n$. But (1) holds for all $V \in \mathfrak{sl}_n(\mathbb{K})$ by assumption, and for all $V \in \mathbb{K} \cdot I_n$ by inspection. Thus, (1) is proven.

Now, consider the map $r : \mathbf{M}_n(\mathbb{K}) \to \mathbf{M}_n(\mathbb{K})$ which maps every V to UVU^{-1}. This r is well-defined due to (1), and a $\mathbb{K}$-algebra isomorphism as can be easily seen; hence, r is a $\mathbb{K}$-algebra automorphism of $\mathbf{M}_n(\mathbb{K})$. But by Skolem-Noether theorem all $\mathbb{K}$-algebra automorphisms of $\mathbf{M}_n(\mathbb{K})$ are inner. Hence, r is inner, so there exists some $P \in \mathrm{GL}(n, \mathbb{K})$ such that every $V \in \mathbf{M}_n(\mathbb{K})$ satisfies $r(V) = PVP^{-1}$. So every $V \in \mathbf{M}_n(\mathbb{K})$ satisfies
$PVP^{-1} = r(V) = UVU^{-1}$.

This can be easily rewritten as $U^{-1}PV\left(U^{-1}P\right)^{-1} = V$. In other words, we have $U^{-1}PV = VU^{-1}P$. Since this holds for all $V \in \mathbf{M}_n(\mathbb{K})$, it must also hold for all $V \in \mathbf{M}_n(\mathbb{L})$ (because it is a linear equation in V, so it is enough to check it on an $\mathbb{L}$-basis of $\mathbf{M}_n(\mathbb{L})$, but such a basis can be chosen to lie in $\mathbf{M}_n(\mathbb{K})$). In other words, the matrix $U^{-1}P$ lies in the center of $\mathbf{M}_n(\mathbb{L})$. But this center is known to be $\mathbb{L} \cdot I_n$. Thus, we obtain $U^{-1}P \in \mathbb{L} \cdot I_n$. In other words, there exists some $\lambda \in \mathbb{L}$ such that $U^{-1}P = \lambda I_n$. This λ is nonzero (else, P would be zero, contradicting $P \in \mathrm{GL}(n, \mathbb{K})$), so this becomes $P = \lambda U$. Hence, we obtain $\lambda U = P \in \mathrm{GL}(n, \mathbb{K})$, proving the lemma.

□

Now we prove Theorem 5.1. That (1) implies (2) is obvious. We consider the statements,

(3) $\mathrm{Ad}(\Delta)$ is definable over $\mathbb{A}$, and
(4) $\mathrm{Ad}(\Gamma)$ is definable over $\mathbb{A}$.

By Proposition 4.3, (2) implies (3). By 4.4, (3) implies (4). We will use the statment (4) to complete the proof that (2) implies (1). Assume (2). Then there exists an invertable linear map g from $\mathbb{F}^{n+1}$ to $\mathbb{F}^{n+1}$ such that $\mathbb{K}^{n+1}$ is an invariant $\mathbb{K}$-form under $g\Delta g^{-1}$. Then the set

$$\mathfrak{sl}_{n+1}(\mathbb{K}) = \{x \in \mathfrak{sl}_{n+1}(\mathbb{F}) \,|\, x \in \mathrm{End}\mathbb{K}^{n+1}\}$$

is a $\mathbb{K}$-form of $\mathfrak{sl}_{n+1}(\mathbb{F})$ invariant under $\mathrm{Ad}(g\Delta g^{-1})$. By proof of Theorem 3 of [29], $\mathfrak{sl}_{n+1}(\mathbb{K})$ is also invariant under $\mathrm{Ad}(g\Gamma g^{-1})$. For a linear transformation y from $\mathbb{F}^{n+1}$ to $\mathbb{F}^{n+1}$, let $\hat{y}$ denote the matrix for y with respect to the standard basis of $\mathbb{F}^{n+1}$. Then since $\mathfrak{sl}_{n+1}(\mathbb{K})$ is invariant under $\mathrm{Ad}(g\Gamma g^{-1})$, we have that for every $x \in \mathfrak{sl}_{n+1}(\mathbb{K})$, $\hat{g}\hat{\sigma}_i\hat{g}^{-1}x\hat{g}\hat{\sigma}_i\widehat{g^{-1}}$ is in $\mathbf{M}_{n+1}(\mathbb{K})$. By Lemma 5.2, there exists a nonzero $\lambda_i \in \mathbb{F}$ such that $\lambda_i\hat{g}\hat{\sigma}_i\hat{g}^{-1}$ is in $\mathrm{GL}(n+1, \mathbb{K})$ so that $\det(\lambda_i\hat{g}\hat{\sigma}_i\hat{g}^{-1}) = \lambda_i^{n+1}$ and $\mathrm{tr}\lambda_i\hat{g}\hat{\sigma}_i\hat{g}^{-1}$ is in $\mathbb{K}$. If σ_i is of the finite order l_i, we have that $(\lambda_i\hat{g}\hat{\sigma}_i\hat{g}^{-1})^{l_i} = \lambda_i^{l_i}Id$ so that $\lambda_i^{l_i}$ is in $\mathbb{K}$. Now if l_i is relatively prime to $n+1$ or nonzero $\mathrm{tr}\sigma_i$ is in $\mathbb{K}$, then we have that λ_i is in $\mathbb{K}$. Hence $\hat{g}\hat{\sigma}_i\widehat{g^{-1}}$ is in $\mathrm{SL}^{\pm}(n+1, \mathbb{K})$ for every i. This shows that Γ is definable over $\mathbb{K}$. Now assume (4). Then by Proposition 4.3 (1) is obtained. □

5.2. **Result for groups generated by reflections.**

Theorem 5.3. *Suppose $n \geq 2$ is an integer. and let $\mathbb{F}$ be an algebraically closed field of characteristic 0. Let $\Gamma \subset \mathrm{SL}^{\pm}(n+1, \mathbb{F})$ be a Zariski dense subgroup generated by finitely many reflections. Let Δ be any subgroup of finite index in Γ. Then the following properties of an integrally closed Noetherian ring $\mathbb{A}$ in $\mathbb{F}$ are equivalent.*

(1) *Γ is definable over $\mathbb{A}$.*
(1′) *$\mathbb{A}$ contains all the simple cyclic products of the Cartan matrix of Γ.*
(2) *Δ is definable over $\mathbb{A}$.*

Proof. Since Γ is Zariski dense in $\mathrm{SL}^{\pm}(n+1, \mathbb{F})$, it is irreducible. By Propositions 4.6 and 4.5 (1) and (1′) are equivalent. Since the trace of a reflection element in $\mathrm{SL}^{\pm}(n+1, \mathbb{F})$ is $n-1$, the equivalence of (1) and (2) is the corollary of Theorem 5.1. □

Now let Γ be a discrete subgroup of $\mathrm{SL}^{\pm}(n+1, \mathbb{R})$ that divides a strictly convex open domain Ω in $\mathbb{S}^n$. By Theorem 2.1, the real Zariski closure $\overline{\Gamma}$ is either $\mathrm{O}^{+}(1, n)$ or $\mathrm{SO}^{+}(1, n)$ iff Ω is an $\mathbb{H}^n$, otherwise $\overline{\Gamma}$ is either $\mathrm{SL}^{\pm}(n+1, \mathbb{R})$ or $\mathrm{SL}(n+1, \mathbb{R})$. Hence if Γ is a polyhedral reflection group which divides a strictly convex open Ω in $\mathbb{S}^n$, then the real Zariski closure $\overline{\Gamma}$ is either $\mathrm{O}^{+}(1, n)$ or $\mathrm{SL}^{\pm}(n+1, \mathbb{R})$. Now Thereom 1.1 is a corollary to Theorem 5.3. □

5.3. **Result for finitely generated groups.** In this section, we state and prove corollaries of Theorem 5.3 in the case when the generators are the special type of the following well-known lemma. For example see [16].

Lemma 5.4. *Let A_θ be the matrix$\begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix}$. An element g in $\mathrm{SL}^{\pm}(n+1, \mathbb{R})$ has finite order if and only if g is similar to*

$$A_{\theta_1}^{[d_1]} \oplus \cdots \oplus A_{\theta_r}^{[d_r]} \oplus I_{k_1} \oplus -I_{k_2},$$

where k_1, $k_2 \geq 0$, $r \geq 0$, $d_1, \cdots, d_r \geq 1$, $0 < \theta_1 < \cdots < \theta_r < \pi$, each θ_i is a rational multiple of 2π, and $k_1 + k_2 + 2(d_1 + \cdots d_r) = n+1$. Wrting $\theta_i = 2\pi a_i/b_i$ with a_i/b_i in lowest terms, the order of such a g is $lcm\{2, b_1 \cdots, b_r\}$ or $lcm\{b_1 \cdots, b_r\}$ according as $k_2 > 0$ or $k_2 = 0$ respectively.

Corollary 5.5. *Let $n \geq 4$. Let $\Gamma \subset \mathrm{SL}(n+1, \mathbb{R})$ be a subgroup generated by finitely many elements. Suppose that we have a presentation of Γ so that every generator g_i of Γ associated to the presentation is of finite order b_i and g_i is similar to the type $A_{\theta_1} \oplus I_{k_1}$ of Lemma 5.4 . Let $\mathbb{A}$ be an integrally closed Noetherian ring in $\mathbb{R}$. Suppose that the real Zariski closure of Γ is $\mathrm{SL}(n+1, \mathbb{R})$. Let Δ be any subgroup of finite index in Γ. Then the following properties of the ring $\mathbb{A}$ are equivalent.*

(1) *Γ is definable over $\mathbb{A}$.*
(2) *Δ is definable over $\mathbb{A}$.*

Proof. By assumption we have that g_i is similar to $A_{\theta_1} \oplus I_{k_1}$ where $\theta_1 = \frac{2a_i\pi}{b_i}$ with a_i a positive integer relatively prime to b_i. Since some power of $A_{\theta_1} \oplus I_{k_1}$ is $A_{\frac{2\pi}{b_i}} \oplus I_{k_1}$ and vice versa, we can assume that $\theta_1 = \frac{2\pi}{b_i}$. We have that $\mathrm{trAd}g_i = (\mathrm{tr}g_i)^2 - 1 = 4\cos^2\frac{2\pi}{b_i} + 4k_1\cos\frac{2\pi}{b_i} + k_1^2 - 1$. Hence $\mathbb{Q}(\mathrm{tr}g_i)$ is at most degree 2 extension over $\mathbb{Q}(\mathrm{trAd}g_i)$. We claim that $\mathbb{Q}(\mathrm{trAd}g_i) = \mathbb{Q}(\mathrm{tr}g_i)$. The claim implie that $\mathrm{tr}g_i$ is in $\mathbb{K}$ since $\mathbb{Q}(\mathrm{trAd}\Gamma)$ is the smallest field of definition of $\mathrm{Ad}\Gamma$ by [29]. Note that $\mathrm{tr}g_i$ is never zero. Hence the theorem will follow from Theorem 5.1. Now we prove the claim.

If $\mathbb{Q}(\mathrm{trAd}g_i) \neq \mathbb{Q}(\mathrm{tr}g_i)$, then $\mathbb{Q}(\mathrm{tr}g_i)$ is a degree 2 extension of $\mathbb{Q}(\mathrm{trAd}g_i)$. Thus there exists a non trivial automorphism ρ of $\mathbb{Q}(\mathrm{tr}g_i)$ which is identity restricted on $\mathbb{Q}(\mathrm{trAd}g_i)$. Namely, there exists a nontrivial $\rho \in \mathrm{Aut}(\mathbb{Q}(\cos\theta_1)/\mathbb{Q}(\cos^2\theta_1 + k_1\cos\theta_1))$. Since $\cos\theta_1$ and $-k_1 - \cos\theta_1$ are two roots of the polynomial

$$x^2 + x - (\cos^2\theta_1 + \cos\theta_1),$$

we have that $\rho(\cos\theta_1) = -k_1 - \cos\theta_1$. But since $\mathbb{Q}(\cos\theta_1) = \mathbb{Q}(\zeta_n + \zeta_n^{-1})$ where $\zeta_n = e^{i\theta_1}$, we have that $\rho(\cos\theta_1) = \cos m\theta_1$ for some positive integer m. Since

$$|\cos m\theta_1| \leq 1 < |-k_1 - \cos\theta_1|$$

we have a contradiction and this proves the claim. □

For subgroups of $\mathrm{SL}(3,\mathbb{R})$ generated by finitely many elements of finite order, every generator is similar to $A_{\theta_1} \oplus I_{k_1}$ where $k_1 = 1$. Hence the same result holds as Corollary 5.5 if there is no generator of order 3. Since the element of order 3 has trace 0, we cannot apply Theorem 5.3 in such cases.

Corollary 5.6. *Let $\Gamma \subset \mathrm{SL}(3,\mathbb{R})$ be a subgroup generated by finitely many elements. Suppose that we have a presentation of Γ so that every generator g_i of Γ associated to the presentation is of finite order and no g_i is of order 3. Let $\mathbb{A}$ be an integrally closed Noetherian ring in $\mathbb{R}$. Suppose that the real Zariski closure of Γ is $\mathrm{SL}(3,\mathbb{R})$. Let Δ be any subgroup of finite index in Γ. Then the following properties of the ring $\mathbb{A}$ are equivalent.*

(1) *Γ is definable over $\mathbb{A}$.*
(2) *Δ is definable over $\mathbb{A}$.*

For subgroups of $\mathrm{SL}(4,\mathbb{R})$ generated by finitely many elements of finite order which is similar to $A_{\theta_1} \oplus I_{k_1}$ where $k_1 = 2$, the same result holds as Corollary 5.5 if there is no generator of order 2. Since the element of order 2 has trace 0, we cannot apply Theorem 5.3 in such cases.

Corollary 5.7. *Let $\Gamma \subset \mathrm{SL}(4,\mathbb{R})$ be a subgroup generated by finitely many elements. Suppose that we have a presentation of Γ so that every generator g_i of Γ associated to the presentation is of finite order and no g_i is of order 2. Suppose further that g_i is similar to the type $A_{\theta_1} \oplus I_2$ of lemma 5.4. Let $\mathbb{A}$ be an integrally closed Noetherian ring in $\mathbb{R}$. Suppose that the real Zariski closure of Γ is $\mathrm{SL}(4,\mathbb{R})$. Let Δ be any subgroup of finite index in Γ. Then the following properties of the ring $\mathbb{A}$ are equivalent.*

(1) *Γ is definable over $\mathbb{A}$.*
(2) *Δ is definable over $\mathbb{A}$.*

6. Examples

6.1. **A preliminary lemma.** In this section, for some 2- , 3- and 4-dimensional Coxeter orbifold $\hat{P}$, we find all or some conjugacy classes of irreducible convex dividing projective reflection groups that are definable over $\mathbb{Z}$ or some other ring of integers. The following lemma of Vinberg [28] will be useful in this section.

Lemma 6.1. *Let $\hat{P}$ be a compact n-dimensional Coxeter orbifold. Suppose that Γ_1 and Γ_2 are convex projective reflection groups and that Ω_1 and Ω_2 are irreducible properly convex open domains that the groups divide respectively. Then the following statements are equivalent.*

(1) Ω_1/Γ_1 *and* Ω_2/Γ_2 *are projectively diffeomorphic.*
(2) *A Cartan matrix* A *of* Γ_1 *and a Cartan matrix* B *of* Γ_2 *are equivalent. Namely* $A = DBD^{-1}$ *for a diagonal matrix* D *having positive diagonal elements.*
(3) *The cyclic products of* A *and the cyclic products of* B *are identical.*

Proof. The equivalence of (2) and (3) follows from Proposition 16 of [28]. Since Γ_1 and Γ_2 divide irreducible properly convex open domains, Vey irreducibility theorem (Theorem 5.1 of [6]) implies that Γ_1 and Γ_2 are irreducible. Thus by Corollary to Proposition 19 of [28], the characteristics of Γ_1 and Γ_2 defined in [28] are determined just by equivalence classes of Cartan matrices. By Theorem 5 of [28], (2) and (1) are equivalent. □

Suppose that Q is a compact hyperbolic orbifold. We recall that every finite index subgroup of fundamental group $\pi_1(Q)$ has a trivial center. Let Γ be a group isomorphic to $\pi_1(Q)$ that divides a properly convex open domain Ω in $\mathbb{S}^n$. Then Corollary 2.13 of Benoist [4] implies that such a group Γ in $\mathrm{SL}^{\pm}(n+1,\mathbb{R})$ is strongly irreducible so that Ω has to be irreducible. By Theorem 1.1 of [2], Ω moreover has to be strictly convex. Hence for a compact hyperbolic Coxeter orbifold $\hat{P}$, we can apply Theorem 1.1 to find out whether an irreducible convex dividing reflection group isomorphic to $\pi_1(\hat{P})$ that is definable over $\mathbb{Z}$ exists or not.

6.2. **Orbifolds based on simplices.** The (p, q, r)- *triangle* is defined to be the triangle with dihedral angles $\frac{\pi}{p}$, $\frac{\pi}{q}$, $\frac{\pi}{r}$. The corresponding Coxeter orbifold is said to be a (p, q, r)-*t*riangular Coxeter orbifold. Let $\hat{P}$ be the (p, q, r)-triangular Coxeter orbifold. It is an elementary fact that $\hat{P}$ is hyperbolic if and only if $\frac{1}{p}+\frac{1}{q}+\frac{1}{r}$ is less than 1. Suppose that $\frac{1}{p}+\frac{1}{q}+\frac{1}{r} < 1$. By condition (2) of Proposition 3.1, we can conclude that for the irreducible convex dividing reflection group isomorphic to $\pi_1(\hat{P})$ to be definable over $\mathbb{Z}$, (p,q,r) must be a triple of elements in $\{2, 3, 4, 6\}$.

Proposition 6.2. *Suppose that* $\frac{1}{p}+\frac{1}{q}+\frac{1}{r} < 1$. *Let* $\hat{P}$ *be a* (p, q, r)*-triangular Coxeter orbifold. The following list shows the number of conjugacy classes of irreducible convex dividing reflection groups isomorphic to* $\pi_1(\hat{P})$ *in* $\mathrm{SL}^{\pm}(3,\mathbb{R})$ *that are definable over* $\mathbb{Z}$*.*

triple	N	triple	N	triple	N
$(2, 4, 6)$	1	$(3, 4, 4)$	3	$(4, 4, 6)$	6
$(3, 3, 4)$	2	$(3, 4, 6)$	4	$(4, 6, 6)$	5
$(3, 3, 6)$	2	$(4, 4, 4)$	4	$(6, 6, 6)$	4
$(2, 6, 6)$	1	$(3, 6, 6)$	3		

Here N *is the number of conjugacy classes of irreducible convex dividing reflection groups isomorphic to* $\pi_1\hat{P}$ *definable over* $\mathbb{Z}$*.*

Proof. We give a proof when $(p, q, r) = (3, 3, 4)$. The other cases are completely analogous. Let

$$\begin{pmatrix} 2 & c_{12} & c_{13} \\ c_{21} & 2 & c_{23} \\ c_{31} & c_{32} & 2 \end{pmatrix}$$

be the Cartan matrix of reflection group associated with $\hat{P}$. By condition (2) of Proposition 3.1, we can choose reflection generators σ_1, σ_2, and σ_3 so that the values of $c_{12}c_{21}$, $c_{13}c_{31}$, and $c_{23}c_{32}$ are fixed to be 1, 1, and 2 respectively. Then since the product of $c_{12}c_{21}$, $c_{13}c_{31}$, and $c_{23}c_{32}$ are equal to the product of $c_{12}c_{23}c_{31}$ and $c_{13}c_{32}c_{21}$), we can have only two possible different integer tuples of simple cyclic products $(c_{12}c_{21}, c_{13}c_{31}, c_{23}c_{32}, c_{12}c_{23}c_{31}, c_{13}c_{32}c_{21})$:$(1,1,2,1,2)$, $(1,1,2,2,1)$. This is equivalent to say that we can have two different equivalent classes of Cartan matrices whose cyclic products are integers. Hence we have two conjugacy classes of irreducible convex dividing reflection groups isomorphic to $\hat{P}$ that are definable over $\mathbb{Z}$.

□

Remark 6.3. The fundamental group $\pi_1(\hat{P})$ has the following Coxeter group presentation,

$$\langle s_1, s_2, s_3 | s_1^2 = s_2^2 = s_3^2 = (s_1s_2)^3 = (s_2s_3)^3 = (s_3s_1)^4 = 1\rangle.$$

Let H be the group which has the following presentation,

$$\langle s_1, s_2 | (s_1^3 = s_2^3 = (s_1s_2)^4 = 1\rangle.$$

Theorem 1.1 implies that there are only two conjugacy classes of irreducible convex dividing linear group Δ which satisfies the following properties.

- Δ is isomorphic to H.
- Δ is a subgroup of irreducible convex dividing (3,3,4) triangle reflection group.
- Δ is definable over $\mathbb{Z}$.

We can compare this to the known result by Long, Reid, and Thistlethwaite ([18]); there are infinitely many conjugacy classes of irreducible convex dividing linear group Δ which satisfies the following properties.

- Δ is isomorphic to H..
- Δ is definable over $\mathbb{Z}$.

Sometimes it is possible to find a 1-parameter family of Cartan matrices such that the set of associated irreducible dividing reflection groups contain all which are definable over $\mathbb{Z}$. For example, let

$$\begin{pmatrix} 2 & -3/2 & -1 \\ -2 & 2 & -2t \\ -2 & -1/2t & 2 \end{pmatrix}$$

be the matrix with $t > 0$. Then this 1-parameter family of matrices satisfies the condition to be a cartan matrix for $(3,4,6)$ triangular irreducible dividing reflection group. This family contains all of four groups which are definable over $\mathbb{Z}$; these are the cases when t is equal to $\frac{1}{6}$, $\frac{1}{3}$, $\frac{1}{2}$, and 1. The Figures 1-4 are images of the convex domains which the four groups divide. These figures were drawn from TrianglegroupProj2, the Mathematica notebook written by the second author([12]).

Proposition 6.4. *Suppose that* $\frac{1}{2} + \frac{1}{q} + \frac{1}{r} < 1$. *Let* $\hat{P}$ *be a* $(2,q,r)$*-triangular Coxeter orbifold. Then there exists a unique conjugacy class of irreducible convex dividing reflection group isomorphic to* $\pi_1(\hat{P})$ *in* $\mathrm{SL}^{\pm}(3,\mathbb{R})$ *and its smallest ring of definition is* $\mathbb{Z}[4\cos^2(\frac{\pi}{q}), 4\cos^2(\frac{\pi}{r})]$.



FIGURE 1. the $(3,4,6)$ triangular reflection group in the case when $t = \frac{1}{6}$



FIGURE 2. the $(3,4,6)$ triangular reflection group in the case when $t = \frac{1}{3}$



FIGURE 3. the $(3,4,6)$ triangular reflection group in the case when $t = \frac{1}{2}$



FIGURE 4. the $(3,4,6)$ triangular reflection group in the case when $t = 1$

Proof. Let

$$\begin{pmatrix} 2 & c_{12} & c_{13} \\ c_{21} & 2 & c_{23} \\ c_{31} & c_{32} & 2 \end{pmatrix}$$

be the Cartan matrix of a irreducible convex dividing reflection group which satisfies the equations $c_{12}c_{21} = 4\cos^2\frac{\pi}{2}$, $c_{13}c_{31} = 4\cos^2\frac{\pi}{q}$ and $c_{23}c_{32} = 4\cos^2\frac{\pi}{r}$. Then the tuple of simple cyclic product $(c_{12}c_{21},\ c_{13}c_{31},\ c_{23}c_{32},\ c_{12}c_{23}c_{31},\ c_{13}c_{32}c_{21})$ has to be $(0, 4\cos^2(\frac{\pi}{q}), 4\cos^2(\frac{\pi}{r}), 0, 0)$. This determines the irreducible convex dividing reflection group up to conjugation. By Lemma 4.5 and 4.6, $\mathbb{Z}[4\cos^2(\frac{\pi}{q}), 4\cos^2(\frac{\pi}{r})]$ is its smallest ring of definition. $\square$

In Section 7.2 of Ratcliffe [25], n-dimensional simplicial Coxeter orbifolds admitting a hyperbolic structure are classified; indeed there is no n-dimensional hyperbolic simplicial orbifold for $n > 4$. For $n = 2$ (Proposition 6.2), we already have classified all the simplicial orbifolds $\hat{P}$s admitting hyperbolic structures such that an irreducible convex dividing reflection group isomorphic to $\pi_1(\hat{P})$ that is definable over $\mathbb{Z}$ exists. For $n = 3$, there are 9 simplicial Coxeter orbifolds admitting hyperbolic structures. Among them, there are only two tetrahedral Coxeter orbifold $\hat{P}$s whose dihedral angles are elements of $\{\frac{\pi}{2}, \frac{\pi}{3}, \frac{\pi}{4}, \frac{\pi}{6}\}$. The condition (2) of Proposition 3.1 and Theorem 1.1 imply that this is a necessary condition for a compact Coxeter orbifold $\hat{P}$ to make irreducible convex dividing reflection groups definable over $\mathbb{Z}$. The two Coxeter tetrahedrons are described in Figure 5. For $n = 4$, there are only 5 simplicial Coxeter orbifolds admitting hyperbolic structures . Among them there are only one Coxeter orbifold $\hat{P}$ whose dihedral angles are elements of $\{\frac{\pi}{2}, \frac{\pi}{3}, \frac{\pi}{4}, \frac{\pi}{6}\}$. The Coxeter graph of this 4-dimensional simplex is described in Figure 6. The following two propositions complete the classification of simplicial orbifolds that admits hyperbolic structure whose projective holonomy groups definable over $\mathbb{Z}$ exists.



FIGURE 5. Two compact Coxeter tetrahedrons definable over $\mathbb{Z}$, $d = 3$ or 4

Proposition 6.5. *Let $\hat{P}$ be one of tetrahedral Coxeter orbifold described in Figure 5. If $d = 3$, then the number of conjugacy classes of the irreducible convex dividing reflection groups isomorphic to $\pi_1(\hat{P})$ in $\mathrm{SL}^{\pm}(4, \mathbb{R})$ that are definable over $\mathbb{Z}$ is 2. If $d = 4$, then the number of conjugacy classes of the irreducible convex dividing reflection groups isomorphic to $\pi_1(\hat{P})$ in $\mathrm{SL}^{\pm}(4, \mathbb{R})$ that are definable over $\mathbb{Z}$ is 3.*

Proof. Let

$$\begin{pmatrix} 2 & c_{12} & c_{13} & c_{14} \\ c_{21} & 2 & c_{23} & c_{24} \\ c_{31} & c_{32} & 2 & c_{34} \\ c_{41} & c_{42} & c_{43} & 2 \end{pmatrix}$$

be the Cartan matrix of the reflection group. By condition (2) of Proposition 3.1, we can choose reflection generators σ_1, σ_2, σ_3, and σ_4 so that the values of $c_{12}c_{21}$, $(c_{13}c_{31}$, c_{14}, c_{41}, c_{23},c_{32}, $c_{24}c_{42}$, $c_{34}c_{43}$ are fixed to be 2, 1, 0, 0, 0, 0, 1, and $4\cos^2\frac{\pi}{d}$ respectively. Now it is easy to check that every simple cyclic product of order 3 is 0, and only nonzero simple cyclic products of order 4 are $c_{12}c_{24}c_{43}c_{31}$ and $c_{13}c_{34}c_{42}c_{21}$.

Finally since the product of $c_{12}c_{21}$, $c_{13}c_{31}$, $c_{24}c_{42}$ and $c_{34}c_{43}$ is equal to the product of $c_{12}c_{24}c_{43}c_{31}$ and $c_{13}c_{34}c_{42}c_{21}$, if $d = 3$, we conclude that only possible integer tuples of nonzero simple cyclic products ($c_{12}c_{21}$, $c_{13}c_{31}$, $c_{24}c_{42}$, $c_{34}c_{43}$, $c_{12}c_{24}c_{43}c_{31}$, $c_{13}c_{34}c_{42}c_{21}$) are (2, 1, 1, 1, 2, 1) and (2, 1, 1, 1, 1, 2). This is equivalent to say that we can have two different equivalence classes of Cartan matrices whose cyclic products are integers. If $d = 4$, we conclude that only possible integer tuples of nonzero simple cyclic products ($c_{12}c_{21}$, $c_{13}c_{31}$, $c_{24}c_{42}$, $c_{34}c_{43}$, $c_{12}c_{24}c_{43}c_{31}$, $c_{13}c_{34}c_{42}c_{21}$) are (2, 1, 1, 2, 4, 1), (2, 1, 1, 2, 1, 4), and (2, 1, 1, 2, 2, 2). This is equivalent to say that we can have three different equivalence classes of Cartan matrices whose cyclic products are integers. □

Proposition 6.6. *Let $\hat{P}$ be the 4-dimensional simplicial Coxeter orbifold whose Coxeter graph is described in Figure 6. Then the number of conjugacy classes of the irreducible convex dividing reflection groups isomorphic to $\pi_1(\hat{P})$ in $\mathrm{SL}^{\pm}(5, \mathbb{R})$ that are definable over $\mathbb{Z}$ is 2.*



FIGURE 6. Coxeter graph of the 4-dimensional simplex definable over $\mathbb{Z}$

Proof. Let

$$\begin{pmatrix} 2 & c_{12} & c_{13} & c_{14} & c_{15} \\ c_{21} & 2 & c_{23} & c_{24} & c_{25} \\ c_{31} & c_{32} & 2 & c_{34} & c_{35} \\ c_{41} & c_{42} & c_{43} & 2 & c_{45} \\ c_{51} & c_{52} & c_{53} & c_{54} & 2 \end{pmatrix}$$

be the Cartan matrix of the reflection group. By condition (2) of Proposition 3.1, we can choose reflection generators σ_1, σ_2, σ_3, σ_4, and σ_5 so that the values of $c_{12}c_{21}$, ($c_{13}c_{31}$, $c_{14}c_{41}$, $c_{15}c_{51}$, $c_{23}c_{32}$, $c_{24}c_{42}$, $c_{25}c_{52}$, $c_{34}c_{43}$, $c_{35}c_{53}$, and $c_{45}c_{54}$ are fixed to be 1, 0, 0, 1, 1, 0, 0, 2, 0, and 1 respectively. Now it is easy to check that every simple cyclic product of order 3 or 4 is 0, and only nonzero simple cyclic products of order 5 are $c_{12}c_{23}c_{34}c_{45}c_{51}$ and $c_{15}c_{54}c_{43}c_{32}c_{21}$.

Finally since the product of $c_{12}c_{21}$, $c_{13}c_{31}$, $c_{24}c_{42}$, $c_{34}c_{43}$ and $c_{45}c_{54}$ is equal to the product of $c_{12}c_{23}c_{34}c_{45}c_{51}$ and $c_{15}c_{54}c_{43}c_{32}c_{21}$, we conclude that only possible integer tuples of nonzero simple cyclic products ($c_{12}c_{21}$, $c_{23}c_{32}$, $c_{34}c_{43}$, $c_{45}c_{54}$, $c_{12}c_{23}c_{34}c_{45}c_{51}$, $c_{15}c_{54}c_{43}c_{32}c_{21}$) are (1, 1, 1, 2, 1, 1, 2) and (1, 1, 1, 2, 1, 2, 1). This is equivalent to say that we can have two different equivalence classes of Cartan matrices whose cyclic products are integers. □

Theorem 6.7. *Let $\hat{P}$ be an n-dimensional simplicial Coxeter orbifold admitting a hyperbolic structure whose Coxeter graph is not a tree. Let Γ be a reflection holonomy group of a properly convex projective structure on $\hat{P}$. Suppose that k is a real number field that is also a field of definition of Γ. Suppose that $k \neq \mathbb{Q}$. Then* *there are infinitely many conjugacy classes of irreducible convex dividing reflection groups isomorphic to* $\pi_1(\hat{P})$ *that are definable over the ring of integers* O_k*.*

Proof. Let

$$\begin{pmatrix} c_{ij} \end{pmatrix}$$

be Cartan matrices which satisfies the condition (2) of Proposition 3.1. By classification of hyperbolic simplicial orbifolds (Figure 7.2.9. of [25]), if the Coeter graph of a hyperbolic simplex is not a tree, then it is a polygon with $n+1$ vertices. Since the Coxeter graph is a polygon, there are two nonzero cyclic products $c_{i_1i_2}c_{i_2i_3}...c_{i_ni_{n+1}}c_{i_{n+1}i_1}$ and $c_{i_1i_{n+1}}c_{i_{n+1}i_n}...c_{i_2i_1}$ of order $n+1$. Observe that the product N of $c_{i_1i_2}c_{i_2i_3}...c_{i_ni_{n+1}}c_{i_{n+1}i_1}$ and $c_{i_1i_{n+1}}c_{i_{n+1}i_n}...c_{i_2i_1}$ is equal to the product of $c_{i_ki_{k+1}}c_{i_{k+1}i_k}$s for $1 \leq k \leq n$ and $c_{i_{n+1}i_1}c_{i_1i_{n+1}}$. Since the Coxeter graph is a polygon with $n+1$ vertices, N has to be an integer ≥ 1.

We can choose the entries other than $c_{i_1i_2}$ and $c_{i_2i_1}$ to be in the smallest ring of definition, which is ,by Proposition 4.5, equal to the ring with unit elements generated by the simple cyclic products. For example, we choose $c_{i_1i_2}$ to be a unit u in O_k and $c_{i_2i_1}$ to be $4u^{-1}\cos^2\frac{\pi}{d_{i_1i_2}}$ where $d_{i_1i_2}$ is the integer determined by the condition (2) of Proposition 3.1. For other entries, if $j < k$ we choose $c_{i_ji_k}$ to be -1, and $c_{i_ji_k}$ to be $-4\cos^2\frac{\pi}{d_{i_ji_k}}$. Then we observe that all the simple cyclic products are elements of O_k. In particular $c_{i_1i_2}c_{i_2i_3}...c_{i_ni_{n+1}}c_{i_{n+1}i_1}$ is equal to $(-1)^n 4u\cos^2\frac{\pi}{d_{i_1i_{n+1}}}$. Since there are infinitely many units in O_k there are infinitely many O_k-tuples of simple cyclic products. □

6.3. **Cubical orbifolds.** The following example illustrates that sometimes knowing Cartan matrices of some irreducible convex dividing reflection holonomy groups is enough to determine the existence of one definable over $\mathbb{Z}$.

Proposition 6.8. *Let* $\hat{P}$ *be the Coxeter orbifold denoted "cu21" in* [14] *that is also described in Figure 7. Then there are at least* 3 *conjugacy classes of irreducible convex dividing reflection holonomy groups in* $\mathrm{SL}^{\pm}(4,\mathbb{R})$ *that are definable over* $\mathbb{Z}$*.*



FIGURE 7. cu21

Proof. Gye-Seon Lee has computed that the dimension of infinitesimal restricted deformation space of convex real projective structures on $\hat{P}$ is 1 in his Mathematica file [17]. This gives a one-parameter family of irreducible convex dividing reflection holonomy groups. Based on Lee's computation we have found in [8] that the corresponding Cartan matrices are

$$\begin{pmatrix} 2 & 0 & 0 & -\frac{2}{2+\sqrt{5}t} & -\frac{2}{2+\sqrt{5}t} & -3+\frac{\sqrt{5}t}{2+2\sqrt{5}t} \\ 0 & 2 & -1 & -\sqrt{6} & 0 & 0 \\ 0 & -1 & 2 & 0 & -\sqrt{6} & 0 \\ -1-\frac{1}{2}\sqrt{5}t & -\sqrt{6} & 0 & 2 & 0 & -\frac{2+\sqrt{5}t}{2+2\sqrt{5}t} \\ -1-\frac{1}{2}\sqrt{5}t & 0 & -\sqrt{6} & 0 & 2 & -\frac{2+\sqrt{5}t}{2+2\sqrt{5}t} \\ -3-\frac{1}{2}\sqrt{5}t & 0 & 0 & -\frac{2(1+\sqrt{5}t)}{2+\sqrt{5}t} & -\frac{2(1+\sqrt{5}t)}{2+\sqrt{5}t} & 2 \end{pmatrix}$$

where t is a real parameter. We have found that in order for every simple cyclic product to be an integer, only possible values of t are 0, $-\frac{4}{5\sqrt{5}}$ and $\frac{4}{\sqrt{5}}$, and these give 3 different integer tuples of simple cyclic products. Hence there are at least 3 conjugacy classes of irreducible convex dividing reflection groups isomorphic to $\pi_1(\hat{P})$ which are definable over $\mathbb{Z}$. $\square$

Remark 6.9. The following equalities give the integral cartan matrices that are equivalent to the parameterized cartan matrix of the cubical reflection group when t is 0, $-\frac{4}{5\sqrt{5}}$ and $\frac{4}{\sqrt{5}}$.

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{2}{\sqrt{6}} & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{2}{\sqrt{6}} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & 0 & 0 & -1 & -1 & -3 \\ 0 & 2 & -1 & -\sqrt{6} & 0 & 0 \\ 0 & -1 & 2 & 0 & -\sqrt{6} & 0 \\ -1 & -\sqrt{6} & 0 & 2 & 0 & -1 \\ -1 & 0 & -\sqrt{6} & 0 & 2 & -1 \\ -3 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}$$

$$\times \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{\sqrt{6}}{2} & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{\sqrt{6}}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 & -1 & -1 & -3 \\ 0 & 2 & -1 & -2 & 0 & 0 \\ 0 & -1 & 2 & 0 & -2 & 0 \\ -1 & -3 & 0 & 2 & 0 & -1 \\ -1 & 0 & -3 & 0 & 2 & -1 \\ -3 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{10}{3\sqrt{6}} & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{10}{3\sqrt{6}} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{5}{3} & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{5}{3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 5 \end{pmatrix} \begin{pmatrix} 2 & 0 & 0 & -\frac{5}{3} & -\frac{5}{3} & -5 \\ 0 & 2 & -1 & -\sqrt{6} & 0 & 0 \\ 0 & -1 & 2 & 0 & -\sqrt{6} & 0 \\ -\frac{3}{5} & -\sqrt{6} & 0 & 2 & 0 & -3 \\ -\frac{3}{5} & 0 & -\sqrt{6} & 0 & 2 & -3 \\ -\frac{13}{5} & 0 & 0 & -\frac{1}{3} & -\frac{1}{3} & 2 \end{pmatrix}$$

$$\times \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{3\sqrt{6}}{10} & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{3\sqrt{6}}{10} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{3}{5} & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{3}{5} & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{5} \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 & -1 & -1 & -1 \\ 0 & 2 & -1 & -2 & 0 & 0 \\ 0 & -1 & 2 & 0 & -2 & 0 \\ -1 & -3 & 0 & 2 & 0 & -1 \\ -1 & 0 & -3 & 0 & 2 & -1 \\ -13 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{2}{3\sqrt{6}} & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{2}{3\sqrt{6}} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{3} & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{3} & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{5} \end{pmatrix} \begin{pmatrix} 2 & 0 & 0 & -\frac{1}{3} & -\frac{1}{3} & -\frac{13}{5} \\ 0 & 2 & -1 & -\sqrt{6} & 0 & 0 \\ 0 & -1 & 2 & 0 & -\sqrt{6} & 0 \\ -3 & -\sqrt{6} & 0 & 2 & 0 & -\frac{3}{5} \\ -3 & 0 & -\sqrt{6} & 0 & 2 & -\frac{3}{5} \\ -5 & 0 & 0 & -\frac{5}{3} & -\frac{5}{3} & 2 \end{pmatrix}$$

$$\times \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{3\sqrt{6}}{2} & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{3\sqrt{6}}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 5 \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 & -1 & -1 & -13 \\ 0 & 2 & -1 & -2 & 0 & 0 \\ 0 & -1 & 2 & 0 & -2 & 0 \\ -1 & -3 & 0 & 2 & 0 & -1 \\ -1 & 0 & -3 & 0 & 2 & -1 \\ -1 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}$$

6.4. **Benoist's Prismatic orbifold.** So far, all the examples of irreducible convex dividing reflection holonomy groups we have considered were ones that divide strictly convex domains. Our final example illustrates that we can apply our main theorem (Theorem 5.3) even when the domain which an irreducible convex dividing reflection holonomy group divides is not strictly convex.

Proposition 6.10. *Suppose that d is 3 or 4. Let $\hat{P}$ be a triangular prism with dihedral angles described in Figure 8. Then the number of irreducible convex dividing reflection holonomy groups in $\mathrm{SL}^{\pm}(4,\mathbb{R})$ which are definable over $\mathbb{Z}$ is 0.*



Figure 8. Benoist's prism

Proof. This example is from Benoist [5]. Benoist proved that there exists a family of Zariski dense irreducible convex dividing reflection groups in $\mathrm{SL}^{\pm}(4,\mathbb{R})$ whose cartan matrices are

$$\begin{pmatrix} 2 & -1 & -1 & 0 & 0 \\ -1 & 2 & -t & 0 & 0 \\ -1 & -t^{-1} & 2 & \frac{\mu(1-t)}{t} & 0 \\ 0 & 0 & 1-t & 2 & -2 \\ 0 & 0 & 0 & -\nu & 2 \end{pmatrix}$$

where $t > 1$, $\mu = \frac{4t}{(t-1)^2}\cos^2(\frac{\pi}{d})$, and $\nu = 2+3\mu$. He further proved that the properly convex open domain Ω they divide is not strictly convex. By Marquis [21], indeed these and the cartan matrices with same parameter t where $0 < t < 1$ gives Cartan matrices of all the irreducible convex dividing reflection groups up to equivalence. (See also Theorem 1.1 of [3].) We observe that $-t$ and $-t^{-1}$ are both the values of a simple cyclic product. Since it is impossible for both $-t$ and $-t^{-1}$ to be integers, the Theorem 5.3 implies that there is no reflection group isomorphic to $\pi_1(\hat{P})$ which is definable over $\mathbb{Z}$.

□

References

[1] Y. Benoist, *Automorphismes des cones convexes*, Invent. Math. **141** , no. 1 149–193 (2000)
[2] Y. Benoist, *Convexes divisibles I*, Tata Institute of Fundamental Research. Stud. Math. **17**, 339–374 (2004)
[3] Y. Benoist, *Convexes divisibles II*, Duke Math. J. **120**, 97–120 (2003)
[4] Y. Benoist *Convexes divisibles III*, Ann. Scient. Ec. Norm. Sup. **38**, 793–832 (2005)
[5] Y. Benoist, *Convexes divisibles IV*, Invent. math. **164**, 249–278 (2006)
[6] Y. Benoist, *A servey on divisible convex sets*, Available from http://www.math.u-psud.fr/~benoist
[7] A. Borel, *Linear Algebraic Groups*, 2nd ed., Graduate Texts in Mathematics. **126**, Springer-Verlag, Berlin, New York, (1991)

[8] K. Choi, *cu21integral.nb, Mathematica file*, available at http://mathsci.kaist.ac.kr/~manifold

[9] S. Choi, *Geometric structures on orbifolds and holonomy representations*, Geom. Dedicata **104** , 161–199 (2004)

[10] S. Choi, *The deformation spaces of projective structures on 3-dimensional Coxeter orbifolds*, Geom. Dedicata **119**, 69–90 (2006)

[11] S. Choi, *Geometric structures on 2-orbifolds: exploration of discrete symmetry*, MSJ Memoirs. vol.27, Math Soc. of Japan (2012)

[12] S. Choi, *TrianglegroupProj2, Mathematica file*, available at http://mathsci.kaist.ac.kr/~manifold

[13] M. Davis, *The geometry and topology of Coxeter groups*, London mathematical society monograph, vol 32, Princeton Univ. Press, Princeton, NJ (2008)

[14] S. Choi, G. Lee, and C. Hodgson, *Projective deformations of hyperbolic Coxeter 3-orbifolds*, Geom. Dedicata (2011), published online.

[15] D. Cooper and K. Delp *The marked length spectrum of a projective manifold or orbifold.* Proc. American Math. Soc. **138**, no. 9, 3361–3376 (2010)

[16] R. Koo, *A Classification of Matrices of Finite Order over C, R and Q* Mathematics Magazine, vol. 76, No. 2, 143–148 (2003)

[17] G-S. Lee, *cu21.nb, Mathematica file*, Available at http://mathsci.kaist.ac.kr/~manifold/cudo.zip

[18] D. Long, A. Reid, and M. Thistlethwaite *Zariski dense surface subgroups in SL(3,Z)*, Geometry and Topology **15** , 1–9. (2011)

[19] D. B. McReynolds, *Arithmetic lattices in SU(n,1).* preprint, Available at http://www.ma.utexas.edu/users/dmcreyn/ComplexArithmeticI.pdf (2006)

[20] C. Maclachlan and A. Reid, *The arithmetic of hyperbolic 3-manifolds.* Graduate Texts in Mathematics. **219**, Springer-Verlag, New York (2003)

[21] L. Marquis, *Espace des modules de certains polydres projectifs miroirs* Geom. Dedicata **147**, 47–86 (2010)

[22] J.S. Milne, *Algebraic Groups, Lie Groups, and their Arithmetic Subgroups.* ,Available at www.jmilne.org/math/ (2011)

[23] A. Onishchik and E. Vinberg *Lie Groups and Algebraic Groups.* Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, Translated from the Russian and with a preface by D. A. Leites. (1990)

[24] W. Thurston, *Geometry and topology of 3-manifolds.* Lecture notes. Princeton University (1979) Available at http://www.msri.org/publications/books/gt3m/

[25] J. Ratcliffe, *Foundations of Hyperbolic Manifolds.* 2nd ed., Graduate Texts in Mathematics. **149**, Springer-Verlag, New York (2006)

[26] E. Vinberg and V. Kac *Quasi-homogeneous cones*, Math. Zametki **1**, 347–354 (1967). English translation, Math. Notes **1**, 231–235 (1967)

[27] E. Vinberg, *Discrete groups generated by reflections in Lobacevski spaces.* Mat. Sb. **72** , 471–488 (1967). English translation, Math. USSR-Sb. **1**, 429–444 (1967)

[28] E. Vinberg, *Discrete linear Groups generated by reflections.* Izv. Akad. Nauk SSSR **35** 1072–1112 (1971) English translation, Math. USSR, IZv. **5**, 10831119 (1971)

[29] E. Vinberg, *Rings of definition of dense subgroups of semisimple linear groups.* Math. USSR Izvestija **5** , 45–55 (1971). English traslation, Math. USSR-IZv. **5**(1), 45–55 (1971)

[30] B. Yahaghi, *On irreducible semigroups of matrices with traces in a subfield.* Linear Algebra Appl. **383**, 17–28 (2004)

Department of Mathematical Sciences, KAIST, Daejeon 305-701, Republic of Korea
E-mail address: `kchoi1982@kaist.ac.kr`

Department of Mathematical Sciences, KAIST, Daejeon 305-701, Republic of Korea
E-mail address: `shchoi@gmail.com`